

HOPLOCERCIDAE

- 1989 Hoplocercidae Frost and Etheridge, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 81: 30.
Type genus: Hoplocercus Fitzinger, 1843.
- 1996 Hoplocercinae—Alifanov, Palaeontol. J., 3: 115.
- 1997 Hoplocercinae—Macey, Larson, Ananjeva, and Papenfuss, J. Mol. Evol., 44: 673
- 1998 Hoplocercinae—Schulte, Macey, Larson, and Papenfuss, Mol. Phylogenetic Evol., 10: 374.
- 2001 Hoplocercidae—Frost, Etheridge, Janies, and Titus, Am. Mus. Novit., 3343: 13.
- 2003 Hoplocercidae—Wiens and Etheridge, Herpetologica, 59: 375.
- 2011 Hoplocercinae—Torres-Carvajal, Etheridge, and de Queiroz, Zootaxa, 2754: 1.
Range: Panamá, Colombia, Amazonian Ecuador and adjacent Peru.
Comment: The most recent revision and taxonomic review is by Torres-Carvajal, Etheridge, and de Queiroz, Zootaxa, 2754: 1–44.

Enyaliooides Boulenger

- 1885 *Enyaliooides* Boulenger, Cat. Liz. Brit. Mus. Nat. Hist., 2: 112. Type species: *Enyalius heterolepis* Bocourt, 1874, by subsequent designation of Burt and Burt, Trans. Acad. Sci. St. Louis, 28 : 13.
- 1969 *Enyaliooides*—Etheridge, Bull. Brit. Mus. (Nat. Hist.), Zool., 18: 256; Fig. 11b.
- 2011 *Enyaliooides*—Torres-Carvajal, Etheridge, and de Queiroz, Zootaxa, 2754: 11.
Range: From Panamá southward along the Pacific slopes of western South America through Colombia and Ecuador, and the upper Amazonian Basin of Colombia, Ecuador, Peru, northwestern Bolivia, and western Brazil.
Comment: Torres-Carvajal, Etheridge, and de Queiroz, Zootaxa, 2754: 11–12, discussed the evidentiary ambiguity for monophyly of this taxon.

Enyaliooides altotambo Torres-Carvajal, Venegas, and de Queiroz

- 2015 *Enyaliooides anisolepis* Torres-Carvajal, Venegas, and de Queiroz, ZooKeys, 494: 110, Fig. 1. Type locality: "Alto Tambo, 5 km on road to Placer, Bosque Integral Otokiki, 0.90600°N; -78.60600°W (DD), 620 m, Provincia Esmeraldas, Ecuador". Holotype: Pontif. Univ. Catol. Ecuador (QCAZ) 8073, by original designation.
Range: Known only from two adjacent localities at 620–645 m in the Chocoan rainforests of northwestern Ecuador.
Comment: Confused with *Enyaliooides oshaughnessyi* prior to its naming according to the original publication.

Enyaliooides anisolepis Torres-Carvajal, Venegas, and de Queiroz

- 2015 *Enyaliooides anisolepis* Torres-Carvajal, Venegas, and de Queiroz, ZooKeys, 494: 116, Fig. 5. Type locality: the eastern bank of the Mayo river, 4.5 km ESE Zumba, -4.88605°S, -79.08738°W (DD), 765 m, Provincia Zamora-Chinchipe, Ecuador", Holotype: Pontif. Univ. Catol. Ecuador (QCAZ) 12537, by original designation.
Range: Between 724 and 1742 m elevation on the Amazonian slopes of the Andes in southern Ecuador and northern Peru.
Comment: Most similar to *Enyaliooides heterolepis* according to the original publication.

Enyalioideas azulae Vengas, Torres-Carvajal, Duran, and de Queiroz.

- 2013 *Enyalioideas azulae* Vengas, Torres Carvajal, Duran, and de Querioz, ZooKeys, 277: 72; Figs. 4. Type locality: "Chambirillo close to Check-point 16 of the CAZMO (07°04'8.9"S, 76°00'51.2"W, 1122m), Provincia de Picota, Region, Peru." Holotype CORBIDI 06772.

Range: Known only from the type locality in the montane rainforest of the Rio Huallaga in northeaster Peru at elevation of 1100 m.

Enyalioides binzaedi Vengas, Torres-Carvajal, Duran, and de Queiroz.

- 2013 *Enyalioides binzaedi* Vengas, Torres Carvajal, Duran, and de Querioz, ZooKeys, 277: 79. Figs, 7-9. Type locality: "Chambirillo close to theCheckpoint of the CAZNP ((07°04'8.9"S, 76°00'51.2"W, 1122m), Provincia de Picota, Region, Peru." Holotype CORBIDI 0882.

Range: Known only from the type locality in the montane rainforest of the Rio Huallaga in northeaster Peru at elevation of 1100 m.

Enyalioides cofanorum Duellman

- 1973 *Enyalioides cofanorum* Duellman, Herpetologica, 29: 230; Fig. 2. Type locality: "Santa Cecilia, Provincia Napo [Sucumbíos], Ecuador". Holotype: Univ. Kansas Mus. Nat. Hist. No. 146658, by original designation.

- 2011 *Enyalioides cofanorum*—Torres-Carvajal, Etheridge, and de Queiroz, Zootaxa, 2754: 11.

Range: Amazonian lowlands east of the Andes in Colombia, Ecuador, and Peru, at elevations between 100–1230 m.

Enyalioides heterolepis (Bocourt)

- 1874 *Enyalius heterolepis* Bocourt, Ann. Sci. Nat., Paris, Ser. 5, 19: 1. Type locality: "Veragua". Holotype: Mus. Natl. Hist. Nat. Paris No. 4067.
- 1878 *Enyalius heterolepis*—Bocourt, Miss. Scient. Mex. Amer. Centr., Rech. Zool., 3: Pl. 20, Fig. 14.
- 1885 *Enyalioides heterolepis*—Boulenger, Cat. Liz. Brit. Mus. Nat. Hist., 2: 114.
- 1905 *Enyalioides insulae* Barbour, Bull. Mus. Comp. Zool., 66: 100. Type locality: "Gorgona Island", Colombia. Holotype: Mus. Comp. Zool. No. 6983. Synonymy by Parker, Ann. Mag. Nat. Hist., Ser. 9, 17: 550.
- 1911 *Enyalioides mocquardi* Despax, Miss. Arc. Merid. Equat. Amer. Sud, Rept. Batr. l'Equat., Paris, 9 (fasc. 2): 22. Type locality: "Equateur". Syntypes: Mus. Natl. Hist. Nat. Paris No. 06.226, 06.228. Synonymy by Burt and Burt, Bull. Am. Mus. Nat. Hist., 61: 265.
- 1926 *Enyalioides heterolepis*—Parker, Ann. Mag. Nat. Hist., Ser. 9, 17: 549.
- 1931 *Enyalioides heterolepis*—Burt and Burt, Bull. Am. Mus. Nat. Hist., 61: 265.
- 1970 *Enyalioides heterolepis*—Peters and Donoso-Barros, Bull. U.S. Natl. Mus., 297: 114.
- 2011 *Enyalioides heterolepis*—Torres-Carvajal, Etheridge, and de Queiroz, Zootaxa, 2754: 13.
- Range: Elevations between 0–1000 m in Panama, the western slopes of the Andes and adjacent Pacific lowlands of Colombia and Ecuador, and Gorgona island, Colombia.
- Comment: Lozkat, Hertz, Stadler, Hamad, and Köhler, Herpetol. Rev.: 521, provided a record from Veraguas, Panama, and commented on the distribution within Panama.

Enyaliooides laticeps (Guichenot)

- 1855 *Enyalius laticeps* Guichenot, in Castelnau (ed.), Anim. nouv. l'Amer. Sud, Paris, Part 2, vol. 2(2): 20; Pl. 5, Fig. a–b. Type locality: "il vient Fonteboa, haut Amazone, partie brésilienne". Syntypes: Mus. Natl. Hist. Nat. Paris No. 6821–6822; Mus. Natl. Hist. Nat. Paris No. 6821 considered holotype by Torres-Carvajal, Etheridge, and de Queiroz, Zootaxa, 2754: 19.
- 1855 *Enyalius planiceps* Guichenot, in Castelnau (ed.), Anim. nouv. l'Amer. Sud, Paris, Part 2, vol. 2(2): 21; Pl. 26, Fig. a–b. Type locality: "de Fonteboa, dans le haut Amazone". Syntypes: Mus. Natl. Hist. Nat. Paris No. 6822 and 6822a, according to Torres-Carvajal, Etheridge, and de Queiroz, Zootaxa, 2754: 19. Synonymy by Boulenger, Cat. Liz. Brit. Mus. Nat. Hist., 2: 113.
- 1875 *Enyalius coeurlescens* Cope, J. Acad. Nat. Sci. Philadelphia, Ser. 2, 8: 169. Type locality: none given; but considered to be "the Amazon from Santarem to Peru" by Gans and Vanzolini, Ann. Carnegie Mus., 33: XXX. Holotype: Acad. Nat. Sci. Philadelphia No. 11382, according to XXX. Synonymy by Gans and Vanzolini, Ann. Carnegie Mus., 33: 123.
- 1885 *Enyaliooides laticeps*—Boulenger, Cat. Liz. Brit. Mus. Nat. Hist., 2: 113.
- 1885 *Enyalius caerulescens*—Boulenger, Cat. Liz. Brit. Mus. Nat. Hist., 2: 112. Incorrect subsequent spelling of *Enyalius coeurlescens* Cope, 1875.
- 1897 *Enyaliooides Festae* Peracca, Boll. Mus. Zool. Anat. Comp. Univ. Torino, 12 (300): 3. Type locality: "Valle del Rio Santiago". Holotype: Mus. Zool. Univ. Torino No. 2169. Synonymy by Burt and Burt, Bull. Am. Mus. Nat. Hist., 61: 228. Subspecies status rejected by Avila-Pires, Zool. Verhand., Leiden, 299: 30.
- 1930 *Enyaliooides laticeps laticeps*—Burt and Burt, Proc. U.S. Natl. Mus., 78: 9.
- 1931 *Enyaliooides laticeps festae*—Burt and Burt, Bull. Am. Mus. Nat. Hist., 61: 266.
- 1953 *Enyaliooides laticeps*—Gans and Vanzolini, Ann. Carnegie Mus., 33: 125.
- 1995 *Enyaliooides laticeps*—Avila-Pires, Zool. Verhand., Leiden, 299: 30.
- 2011 *Enyaliooides laticeps*—Torres-Carvajal, Etheridge, and de Queiroz, Zootaxa, 2754: 19.
Range: Throughout the western Amazon basin at elevations between 80–1600 m in Colombia, Ecuador, Peru, and Brazil; possibly extending into Bolivia

Enyaliooides microlepis (O'Shaughnessy)

- 1881 *Enyalius microlepis* O'Shaughnessy, Proc. Zool. Soc. London, 1881 238; Pl. 24, Fig. 2. Type locality: "Sarayacu". Syntypes: Brit. Mus. Nat. Hist. No. 60.6.16.8 (renumbered 1946.8.5.76), 58.7.25.17 (renumbered 1946.8.5.57), and 80.12.8.36–36a (renumbered 1946.8.9.5–6).
- 1885 *Enyaliooides microlepis*—Boulenger, Cat. Liz. Brit. Mus. Nat. Hist., 2: 115.
- 2011 *Enyaliooides microlepis*—Torres-Carvajal, Etheridge, and de Queiroz, Zootaxa, 2754: 22.
Range: East of the Andes in southern Colombia, Ecuador, and northern Peru at elevations between 100–1000 m.

Enyaliooides oshaughnessyi (Boulenger)

- 1881 *Enyalius oshaughnessyi* Boulenger, Proc. Zool. Soc. London, 1881: 246; Pl. 26, Fig. 1, 1a. Type locality: "Ecuador". Holotype: Mus. Roy. Hist. Nat. Brussels, No. 2009.
- 1885 *Enyaliooides oshaughnessyi*—Boulenger, Cat. Liz. Brit. Mus. Nat. Hist., 2: 115.

2011 *Enyaliooides oshaghnessyi*—Torres-Carvajal, Etheridge, and de Queiroz, Zootaxa, 2754: 23.

Range: Western slopes of the Andes and adjacent Pacific lowlands of northern Ecuador and southern Colombia at elevations between 50–1600 m.

Enyaliooides palpebralis (Boulenger)

1883 *Enyalius palpebralis* Boulenger, Proc. Zool. Soc. London, 1883; Pl. 10. Type locality: "Cashiboya, in Eastern Peru". Holotype: Brit. Mus. Nat. Hist. No. 81.5.13.25 (renumbered 1946.8.9.8).

1885 *Enyaliooides palpebralis*—Boulenger, Cat. Liz. Brit. Mus. Nat. Hist., 2: 117.

2004 *Enyaliooides palpebralis*—Reichle, Cortez, Padial, and De la Riva, Bol. Asoc. Herpetol. Esp., 15: 14–16.

2011 *Enyaliooides oshaghnessyi*—Torres-Carvajal, Etheridge, and de Queiroz, Zootaxa, 2754: 25.

Range: Eastern slopes of the Andes and adjacent lowlands in Peru, northern Bolivia, and western Brazil at elevations between 100–1300 m.

Enyaliooides praestabilis (O'Shaughnessy)

1881 *Enyalius praestabilis* O'Shaughnessy, Proc. Zool. Soc. London, 1881: 340; Pl. 25, Fig. 1. Type locality: "Pallatanga and Canelos"; restricted to "Canelos (Provincia Pastaza, 1°34' 60" S, 77° 45' W, 631 m)" by Torres-Carvajal, Etheridge, and de Queiroz, Zootaxa, 2754: 28. Syntypes: Brit. Mus. Nat. Hist. No. 8.12.8.38 (renumbered 1946.8.9.5), 80.12.8.37.

1885 *Enyaliooides praestabilis*—Boulenger, Cat. Liz. Brit. Mus. Nat. Hist., 2: 113.

2011 *Enyaliooides praestabilis*—Torres-Carvajal, Etheridge, and de Queiroz, Zootaxa, 2754: 27.

Range: The Amazon Basin of Ecuador.

Enyaliooides rubrigularis Torres-Carvajal, de Queiroz, and Etheridge

2009 *Enyaliooides rubrigularis* Torres-Carvajal, de Queiroz, and Etheridge, ZooKeys, 27: 61; Fig. 1–4. Type locality "finca de Mesías San Martín (3° 51' 23" S, 78° 51' 53" W, 1154 m), near Piuntza, Provincia Zamora Chinchipe, Ecuador". Holotype: Mus. Zool. Pont. Univ. Catól. Ecuador No. 8483.

Range: Upper basin of the Río Zamora on the eastern slopes of the Andes, and the western slopes of the Corillera del Cóndor, Ecuador.

Enyaliooides rudolfarndti Vegas, Duran, Landauro, and Lujan

2011 *Enyaliooides rudolfarndti* Venegas, Duran, Landauro, and Lujan, Zootaxa, 3109: 40; Fig. 1–3. Type locality "Pan de Azcar trail near the Puesto de Control Huampal in the YCNP (10° 11' 03" S 75° 34' 27" W; WGS 84) at 1050 m.a.s.l. . . . Provincia de Oxapampa, Región de Pasco, Peru". Holotype: Cent. Ornith. Biodivers. No. 07210, by original designation.

Range: Known only from the type locality in the Región de Pasco, upper Amazon Basin of central Peru.

Enyaliooides sophiarothschildae Torres-Carvajal, Venegas, and de Queiroz

2015 *Enyaliooides sophiarothschildei* Torres-Carvajal, Venegas, and de Queiroz, ZooKeys, 494: 121, Fig. 8. Type locality: "Río Lejía in the trail La Cueva-Añazco Pueblo, - 6.83655°S; -77.48603°W (DD), 1700 m, Provincia Mariscal Cáceres, Región San Martín, Perú". Holotype: Div. Herpetol. Centr. Ornithol. Biodivers. (CORBIDI) 647, by original designation.

Range: Elevations of 1600 to 1700 m from two adjacent localities two adjacent localities: 1) the trail to La Cueva-Añasco Pueblo in the drainage of the Lejía river and 2) El Dorado in the drainage of the Blanco river, both tributaries of the Huallabamba river in the northern part of the Huallaga river basin, northeastern slopes of the Cordillera Central in Peru.

Enyaliooides touzeti Torres-Carvajal, Almendáriz, Valencia, Yáñez-Muñoz, and Reyes

2008 *Enyaliooides touzeti* Torres-Carvajal, Almendáriz, Valencia, Yáñez-Muñoz, and Reyes, Pap. Avulsos Zool., São Paulo, 48: 228; Fig 1, 2A–D, 4. Type locality: "Finca La Envidia (3°3'0"S, 79°41'15"W, 433 m) Santa Marta hill, Cantón Ponce Enriquez, Priovincia Azuay, Ecuador". Holotype: Esc. Politech. Nac. Quito No. 10306.

Range: Lowland cloud forests on the western slopes of the Andes in southern Ecuador at elevations of 300–700 m in the provinces of Azuay, Cañar, and El Oro.

Hoplocercus Fitzinger

1843 *Hoplocercus* Fitzinger, Syst. Rept.: 17, 78. Type species: *Hoplocercus spinosus* Fitzinger, 1843, by original designation.

1854 *Pachycercus* Dugès and Braconnier in Duméril, Rev. Mag. Zool., Paris, Ser. 2, 6: 558. Type species: *Pachycercus aculeatus* Dugès and Braconnier, 1854, by monotypy.

1885 *Hoplocercus*—Boulenger, Cat. Liz. Brit. Mus. Nat. Hist., 2: 199.

Range: The state of São Paulo in southeastern Brazil.

Hoplocercus spinosus Fitzinger

1843 *Hoplocercus spinosus* Fitzinger, Syst. Rept.: 17, 78. Type locality: "America. Brasilia". Holotype: presumably originally in Natur. Mus. Wien, not located.

1854 *Pachycercus aculeatus* Dugès and Braconnier in Duméril, Rev. Mag. Zool., Paris, Ser. 2, 6: 561; Pl. 12, Fig. 1-4. Type locality: "Saint-Paul (Brésil)". Holotype: not located. Synonymy by Boulenger, Cat. Liz. Brit. Mus. Nat. Hist., 2: 199.

1885 *Hoplocercus spinosus*—Boulenger, Cat. Liz. Brit. Mus. Nat. Hist., 2: 199.

Range: The state of São Paulo in southeastern Brazil.

Morunasaurus Dunn

1933 *Morunasaurus* Dunn, Occas. Pap. Boston Soc. Nat. Hist., 8: 75. Type species: *Morunasaurus groi* Dunn, 1933, by original designation.

Range: Panamá and Amazonian Ecuador.

Morunasaurus annularis (O'Shaughnessy)

1881 *Hoplocercus annularis* O'Shaughnessy, Proc. Zool. Soc. London, 1881: 244; Pl. 25, Fig. 2. Type locality: "Canelos". Holotype: Brit. Mus. Nat. Hist. No. 80.12.8.54 (renumbered 1946.8.10.35).

1933 *Morunasaurus annularis*—Dunn, Occas. Pap. Boston Soc. Nat. Hist., 8: 76.

Range: Amazonian Ecuador.

Morunasaurus groi Dunn

1933 *Morunasaurus groi* Dunn, Occas. Pap. Boston Soc. Nat. Hist., 8: 76. Type locality: "El Valle de Anton, Panama(8° 36' N, 80° 7' W) about 40 miles south southwest of Barro Colorado Island". Holotype: Mus. Comp. Zool. No. 34875.

Range: Panamá, where known from the type locality and from the Parque Nacional Altos de Campana, and northwestern Colombia.

Morunasaurus peruvianus Köhler

2003 *Morunasaurus peruvianus* Köhler, Senckenb. Biol., 82: 3. Type Locality "vicinity of Kumpin entse, on Río Najem (tributary of the Río Hampami, which is a tributary of the Río Cenapa), Dep. Amazonas, Peru". Holotype: U.S. Natl. Mus. Nat. Hist. No. 316725, by original designation.

Range: Río Cenepa region in northern Peru.

Literature Cited

- Alifanov, V.R. 1996. Lizard families Priscagamidae and Hoplocercidae (Sauria, Iguania): Phylogenetic position and new representatives from the Late Cretaceous of Mongolia. Paleontological Journal 3: 100–118.
- Avila-Pires, T.C.d.S. 1995. Lizards of Brazilian Amazonia (Reptilia: Squamata). Zoologische Verhandelingen. Leiden 299: 1–706.
- Barbour, T. 1905. The Vertebrata of Gorgona Island, Colombia: Reptilia; Amphibia. Bulletin of the Museum of Comparative Zoology 66: 98–102.
- Bocourt, F. 1874. Deux notes sur quelques sauriens de l'Amérique méridionale. Annales des Sciences Naturelles. Zoologie et Biologie Animale. Paris. Serie 5, 19: 1–5.
- Bocourt, F. 1878. Mission Scientifique au Mexique et dans l'Amérique Centrale. Recherches Zoologiques. Troisième Partie, 1 Section. Études sur les reptiles et les batraciens. Livraison 5: 281–360, plates 20, 20D–G, 21A–C. Paris: Ministère de l'Instruction Publique.
- Boulenger, G.A. 1881. Description of a new species of Enyalius in the Brussels Museum. Proceedings of the Zoological Society of London 1881: 246–247.
- Boulenger, G.A. 1883. Description of a new species of lizard of the genus Enyalius. Proceedings of the Zoological Society of London 1883: 46, pl.
- Boulenger, G.A. 1885. Catalogue of the lizards in the British Museum (Natural History). Volume 2. Iguanidae, Xenosauriae, Zonuridae, Anguidae, Annellidae, Helodermatidae, Varanidae, Xantusiidae, Teiidae. London: Printed by Order of the Trustees.
- Burt, C.E., and M.D. Burt. 1930. The South American lizards in the collection of the United States National Museum. Proceedings of the United States National Museum 78: 1–52.
- Burt, C.E., and M.D. Burt. 1931. South American lizards in the collection of the American Museum of Natural History. Bulletin of the American Museum of Natural History 61: 227–395.
- Burt, C.E., and M.D. Burt. 1933. A preliminary check list of the lizards of South America. Transactions of the Academy of Science of Saint Louis 28: i–v, 1–104.

- Cope, E.D. 1875 "1876". On the Batrachia and reptiles collected by Dr. John M. Bransford during the Nicaraguan Canal Survey of 1873. *Journal of the Academy of Natural Sciences of Philadelphia*. Series 2, 8: 155–157.
- Despax, R. 1911. Reptiles et batraciens de l'Équateur recueillis par M. Le Dr. Rivet. , Mission du Service Géographique de l'Armée pour la Mesure d'un Arc de Méridien Équatorial en Amérique du Sud sous le Contrôle Scientifique de l'Académie des Sciences 1899-1906. Tome 9 (Zoologie), Fascicule 2: B.17–B.44, 3 plates. Paris: Gauthier-Villars.
- Duellman, W.E. 1973. Descriptions of new lizards from the upper Amazon Basin. *Herpetologica* 29: 228–231.
- Duméril, A.H.A. 1854. Essai d'application à la classe des reptiles d'une distribution par séries parallèles, et description d'un Saurien, type d'un genre nouveau, dans la famille des Iguaniens (le Pachycerque, Al. Dugès et Sér. Bracconier). *Revue et Magasin de Zoologie Pure et Appliquée*. Serie 2. Paris 6: 544–562.
- Dunn, E.R. 1933. Amphibians and reptiles from El Valle de Anton, Panama. *Occasional Papers of the Boston Society of Natural History* 8: 65–79.
- Etheridge, R.E. 1969. A review of the Iguanid lizard genus *Enyalius*. *Bulletin of the British Museum (Natural History), Zoology* 18: 233–260.
- Fitzinger, L.J.F.J. 1843. *Systema Reptilium. Fasciculus Primus*. Wien: Braumüller et Seidel.
- Frost, D.R., and R.E. Etheridge. 1989. A phylogenetic analysis and taxonomy of iguanian lizards (Reptilia: Squamata). *Miscellaneous Publications. Museum of Natural History, University of Kansas* 81: 1–65.
- Frost, D.R., R.E. Etheridge, D. Janies, and T.A. Titus. 2001. Total evidence, sequence alignment, evolution of polychrotid lizards, and a reclassification of the Iguania (Squamata: Iguania). *American Museum Novitates* 3343: 1–38.
- Gans, C., and P.E. Vanzolini. 1953. Status of the South American iguanid lizard, *Enyalius coeruleascens* Cope. *Annals of the Carnegie Museum* 33: 125–127.
- Guichenot, A. 1855. Reptiles. In F.L.d.L. Castelnau (ed.), *Animaux Nouveaux ou Rares Recueillis Pendant l'Expédition dans les Parties Centrales de l'Amérique du Sud, de Rio de Janeiro à Lima, et de Lima au Para; Exécutée par Ordre du Gouvernement Français Pendant les Années 1843 à 1847, Sous la Dir. de Francis de Castelnau. Part VII. Zoologie*, Volume 2, part 2. Paris: P. Bertrand.
- Köhler, G. 2003. A new species of *Morunasaurus* from Peru (Reptilia, Squamata, Hoplocercidae). *Senckenbergiana Biologica* 82: 235–241.
- Lozkat, S., A. Hertz, L. Stadler, N. Hamad, and G. Köhler. 2010. Noteworthy distribution records of reptiles from western Panama. *Herpetological Review*: 4.
- Macey, J.R., A. Larson, N.B. Ananjeva, and T.J. Papenfuss. 1997. Evolutionary shifts in three major structural features of the mitochondrial genome among iguanian lizards. *Journal of Molecular Evolution* 44: 660–674.
- O'Shaughnessy, A.W.E. 1881. An account of the collection of lizards made by Mr. Buckley in Ecuador. *Proceedings of the Zoological Society of London* 1881: 227–245.
- Parker, H.W. 1926. The reptiles and batrachians of Gorgona Island, Colombia. *Annals and Magazine of Natural History*, Series 9, 17: 549–554.
- Peracca, M.G. 1897. Viaggio del Dott. E. Festa nell'Ecuador e regioni vicini, IV. Rettili. *Bollettino dei Musei di Zoologia e Anatomia Comparata della R. Universita di Torino* 12 (300): 1–20.

- Peters, J.A., and R. Donoso-Barros. 1970. Catalogue of the Neotropical Squamata: Part II. Lizards and amphisbaenians. Bulletin of the United States National Museum 297: i–v, 1–293.
- Reichle, S., C. Cortez, J.M. Padial, and I. De la Riva. 2004. Notes on *Enyalioides palpebralis* (Boulenger, 1883) (Reptilia: Squamata: Hoplocercidae) in Bolivia. Boletin de la Asociacion Herpetologica Espanola 15: 14–16.
- Schulte, J.A., II, J.R. Macey, A. Larson, and T.J. Papenfuss. 1998. Molecular tests of phylogenetic taxonomies: a general procedure and example using four subfamilies of the lizard family Iguanidae. Molecular Phylogenetics and Evolution 10: 367–376.
- Torres-Carvajal, O., A. Almendáriz, J. Valencia, M. Yáñez-Muñoz, and J.P. Reyes. 2008. A new species of *Enyalioides* (Iguanidae: Hoplocercinae) from southwestern Ecuador. Papéis Avulsos de Zoologia. São Paulo 48: 227–235.
- Torres-Carvajal, O., K. de Queiroz, and R. Etheridge. 2009. A new species of iguanid lizard (Hoplocercinae, *Enyalioides*) from southern Ecuador with a key to eastern Ecuadorian *Enyalioides*. ZooKeys 27: 59–71.
- Torres-Carvajal, O., R. Etheridge, and K. de Queiroz. 2011. A systematic revision of Neotropical lizards in the clade *Hoplocercinae* (Squamata: Iguania). Zootaxa 2754: 1–44.
- Torres-Carvajal, O., P.J. Venegas, and K. de Queiroz. 2015. Three new species of woodlizards (Hoplocercinae, *Enyalioides*) from northwestern South America. ZooKeys 494: 107–132.
- Venegas, P.J., V. Duran, C.Z. Landauro, and L. Lujan. 2011. A distinctive new species of wood lizard (Hoplocercinae, *Enyalioides*) from the Yanachaga Chemillen National Park in central Peru. Zootaxa 3109: 39–48.
- Vengas, P.J., O. Torres Carvajal, V. Duran, and K. de Querioz. 2013. Two sympatric new species of woodlizards (Hoplocercinae, *Enyalioides*) from Cordillera Azul National Park in northeastern Peru. ZooKeys 277: 69–90.
- Wiens, J.J., and R.E. Etheridge. 2003. Phylogenetic relationships of hoplocercid lizards: coding and combining meristic, morphometric, and polymorphic data using step matrices. Herpetologica 59: 375–398.